



King's Research Portal

DOI:

[10.4171/QT/131](https://doi.org/10.4171/QT/131)

Document Version

Peer reviewed version

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Etgu, T., & Lekili, Y. (2019). Fukaya categories of plumbings and multiplicative preprojective algebras. *Quantum Topology*, 10(4), 777-813. <https://doi.org/10.4171/QT/131>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

FUKAYA CATEGORIES OF PLUMBINGS AND MULTIPLICATIVE PREPROJECTIVE ALGEBRAS

TOLGA ETGÜ AND YANKI LEKILI

ABSTRACT. Given an arbitrary graph Γ and non-negative integers g_v for each vertex v of Γ , let X_Γ be the Weinstein 4-manifold obtained by plumbing copies of $T^*\Sigma_v$ according to this graph, where Σ_v is a surface of genus g_v . We compute the wrapped Fukaya category of X_Γ (with bulk parameters) using Legendrian surgery extending our previous work [14] where it was assumed that $g_v = 0$ for all v and Γ was a tree. The resulting algebra is recognized as the (derived) multiplicative preprojective algebra (and its higher genus version) defined by Crawley-Boevey and Shaw [8]. Along the way, we find a smaller model for the internal DG-algebra of Ekholm-Ng [12] associated to 1-handles in the Legendrian surgery presentation of Weinstein 4-manifolds which might be of independent interest.

1. INTRODUCTION

In this paper, we continue our study from [14] of computing explicit models for the wrapped Fukaya category $\mathcal{W}(X_\Gamma)$ of the 4-dimensional open symplectic manifold X_Γ , obtained by plumbing cotangent bundles of surfaces according to a graph Γ . Previously in [14], we restricted to case of plumbings of cotangent bundles of 2-spheres and X_Γ was exhibited as a Legendrian surgery along a link $L_\Gamma \subset (S^3, \xi_{std})$ and the Chekanov-Eliashberg DG-algebra \mathcal{B}_Γ of L_Γ was computed explicitly for any tree Γ . It is known that \mathcal{B}_Γ is A_∞ -quasi-isomorphic to the endomorphism algebra of the full subcategory of $\mathcal{W}(X_\Gamma)$ whose objects are the cocores of the 2-handles in the surgery presentation [5] (cf. [11, Thm. 2]). Furthermore, a combination of results of [5] and [1] implies that these cocores split-generate $\mathcal{W}(X_\Gamma)$. A proof of the stronger statement of generation (rather than split-generation) is recently given by Chantraine, Dimitroglou-Rizell, Ghiggini, Golovko [9]. Thus, we take \mathcal{B}_Γ as our primary object of interest.

We improve on the results of [14] in two aspects. Firstly, no restriction is imposed on Γ which was assumed to be a tree in [14] and only plumbings of T^*S^2 's were considered. When Γ is not a tree or cotangent bundles of positive genus surfaces are used, Legendrian surgery description of X_Γ necessarily includes 1-handles, and the computation of \mathcal{B}_Γ is more involved. More precisely, for any connected graph Γ with s vertices and $s + k - 1$ edges and $g = \sum_v g_v$, we shall construct a Legendrian L_Γ in $\#^{k+2g}(S^1 \times S^2, \xi_{std})$ with s components such that X_Γ is obtained by Legendrian surgery on L_Γ . A roadmap for an explicit computation of \mathcal{B}_Γ is provided by the work of Ekholm-Ng [12] which involves a countably generated internal DG-algebra \mathcal{I} associated to each 1-handle in the surgery presentation. We find a simplification by proving that a finitely generated DG-subalgebra of \mathcal{I} is quasi-isomorphic to \mathcal{I} .

Secondly, when Γ is one of the A_n - and D_n -type trees (and conjecturally we also include E_6, E_7, E_8 -types), it was shown in [14] that \mathcal{B}_Γ is quasi-isomorphic to Ginzburg's DG-algebra \mathcal{G}_Γ , away from a small set of exceptional characteristics for the ground field \mathbb{K} . On the other hand, if Γ is a non-Dynkin tree, the explicit computation in [14] shows that \mathcal{B}_Γ is a deformation of \mathcal{G}_Γ . In this case, it is also known that \mathcal{G}_Γ is quasi-isomorphic to $H^0(\mathcal{G}_\Gamma) = \Pi_\Gamma$, the (additive) preprojective algebra attached to Γ (in contrast, \mathcal{G}_Γ has non-trivial cohomology in every non-positive degree, when Γ is a Dynkin tree). It was argued that the associated formal deformation is trivial over a field of characteristic 0 since Π_Γ has no formal deformations as a Calabi-Yau algebra, but the precise determination of quasi-isomorphism type of \mathcal{B}_Γ was not given.

In this paper, we first restrict to the case $g_v = 0$ and show that for any finite graph Γ , \mathcal{B}_Γ is quasi-isomorphic to the derived multiplicative preprojective algebra \mathcal{L}_Γ whose zeroth cohomology is isomorphic to the multiplicative preprojective algebra Λ_Γ defined in [8] by Crawley-Boevey and Shaw with motivation coming from the Deligne-Simpson problem. Next, we extend this to the case where g_v is arbitrary. The resulting DG-algebra is quasi-isomorphic to a positive genus version of the derived multiplicative algebra that appears in [3]. We shall denote this by $\mathcal{L}_{\Gamma, \mathbf{g}}$ and its zeroth cohomology $H^0(\mathcal{L}_{\Gamma, \mathbf{g}})$ by $\Lambda_{\Gamma, \mathbf{g}}$, where $\mathbf{g} = (g_1, g_2, \dots, g_s)$ is an s -tuple of non-negative integers recording the genus g_v associated to vertices of Γ . When \mathbf{g} is omitted, it is understood that $g_v = 0$ for all v .

Before giving the definition of $\Lambda_{\Gamma, \mathbf{g}}$ and $\mathcal{L}_{\Gamma, \mathbf{g}}$, let us first introduce some notation. As in [14], we fix a field \mathbb{K} and the semisimple ring

$$\mathbf{k} := \bigoplus_{v=1}^s \mathbb{K}e_v$$

where $e_v^2 = e_v$ and $e_v e_w = 0$ for $v \neq w$, and $\{1, \dots, s\}$ is the set of vertices of Γ . Here, we shall work over a bigger ring:

$$\mathbf{k}\langle \mathbf{t}^\pm \rangle = \mathbf{k}\langle t_1, t_1^{-1}, t_2, t_2^{-1}, \dots, t_s, t_s^{-1} \rangle / (t_v t_v^{-1} = t_v^{-1} t_v = e_v, t_v^\pm e_w = e_w t_v^\pm = 0 \text{ for } v \neq w)$$

We also fix a spanning tree $T \subset \Gamma$ and orient the edges of Γ . The multiplicative preprojective algebra Λ_Γ is the associative algebra obtained from $\mathbf{k}\langle \mathbf{t}^\pm \rangle$ by adjoining free (non-commuting) generators

$$\begin{aligned} & c_a, c_a^* \text{ for any arrow } a \in \Gamma, \text{ and} \\ & z_a, z_a^{-1} \text{ for any arrow } a \in \Gamma \setminus T \\ & \alpha_{v,i}, \alpha_{v,i}^{-1}, \beta_{v,i}, \beta_{v,i}^{-1} \text{ for any vertex } v \in \{1, \dots, s\} \text{ and } i = 1, 2, \dots, g_v. \end{aligned}$$

(Note that there are indeed 4 generators associated to $a \in \Gamma \setminus T$: $c_a, c_a^*, z_a, z_a^{-1}$.)

The $\mathbf{k}\langle \mathbf{t}^\pm \rangle$ -bimodule structure on Λ_Γ is obtained by extending the scalars from the \mathbf{k} -bimodule structure defined by

$$\begin{aligned} e_w c_a e_v &= c_a, & e_v c_a^* e_w &= c_a^*, \\ e_v z_a^\pm e_v &= z_a^\pm \end{aligned}$$

if and only if $v = s(a)$ and $w = t(a)$ are the source and target vertices of the arrow a , respectively, and \mathbf{k} -bimodule structure on the remaining generators is the obvious one:

$$\begin{aligned} e_v \alpha_{v,i}^\pm e_v &= \alpha_{v,i}^\pm \\ e_v \beta_{v,i}^\pm e_v &= \beta_{v,i}^\pm \end{aligned}$$

In all other cases, the product with an idempotent gives zero. With this understood, the underlying algebra structure of Λ_Γ is the tensor algebra of the \mathbf{k} -bimodule generated by $c_a, c_a^*, z_a^\pm, \alpha_{v,i}^\pm, \beta_{v,i}^\pm, t_v^\pm$.

Note that the variables t_v are taken to be non-central (unless we specialise). The generators of Λ_Γ are subject to the following relations (all multiplications are read from right-to-left):

$$\begin{aligned} z_a z_a^{-1} &= z_a^{-1} z_a = e_v, & \text{for } a \in \Gamma \setminus T \text{ and } v = s(a) \\ e_v + c_a^* c_a &= z_a, & \text{for } a \in \Gamma \setminus T \text{ and } v = s(a) \end{aligned}$$

$$\prod_{t(a)=v} (e_v + c_a c_a^*) = t_v \prod_{i=1}^{g_v} [\alpha_{v,i}, \beta_{v,i}] \prod_{s(a)=v} (e_v + c_a^* c_a), \quad \text{for } v \in \{1, \dots, s\}$$

where the terms in the products are ordered according to a fixed order of all the arrows in Γ , and the empty product is understood to be the relevant idempotent. The notation $[x, y]$ stands for the commutator $xyx^{-1}y^{-1}$.

It is known that $\Lambda_{\Gamma, \mathbf{g}}$ is independent of the orientation and order of the edges of Γ , and the choice of the spanning tree T (cf. [8, Thm. 1.4]). This also follows from our main result by the invariance properties of Chekanov-Eliashberg DG-algebra $\mathcal{B}_{\Gamma, \mathbf{g}}$.

A DG-algebra $\mathcal{L}_{\Gamma, \mathbf{g}}$ such that $H^0(\mathcal{L}_{\Gamma, \mathbf{g}}) \simeq \Lambda_{\Gamma, \mathbf{g}}$ is discussed in [3, Rem.5C & Ex.5.6] (where t_ν are called q_ν which are taken as parameters in \mathbb{C}^\times , see also [27]) under the name (universal) derived multiplicative preprojective algebra. In the current set-up, $\mathcal{L}_{\Gamma, \mathbf{g}}$ can be defined by adding generators ζ_a for $a \in \Gamma \setminus T$ and τ_v for $v \in \{1, \dots, s\}$ with grading -1 , in lieu of the last two relations above, to the set of generators of $\Lambda_{\Gamma, \mathbf{g}}$ (which are assigned grading 0) so that $H^0(\mathcal{L}_{\Gamma, \mathbf{g}}) = \Lambda_{\Gamma, \mathbf{g}}$. Namely, the differentials of these additional generators are

$$\begin{aligned} \partial \zeta_a &= e_v + c_a^* c_a - z_a, & \text{for } a \in \Gamma \setminus T \text{ and } v = s(a) \\ \partial \tau_v &= \prod_{t(a)=v} (e_v + c_a c_a^*) - t_v \prod_{i=1}^{g_v} [\alpha_{v,i}, \beta_{v,i}] \prod_{s(a)=v} (e_v + c_a^* c_a), & \text{for } v \in \{1, \dots, s\} \end{aligned}$$

We can now state our main theorem succinctly as follows:

Theorem 1. *For any graph Γ and \mathbf{g} , there is a quasi-isomorphism between the derived multiplicative preprojective algebra $\mathcal{L}_{\Gamma, \mathbf{g}}$ and the Chekanov-Eliashberg DG-algebra $\mathcal{B}_{\Gamma, \mathbf{g}}$.*

Remark 2. The main theorem is proved by identifying $\mathcal{B}_{\Gamma, \mathbf{g}}$ and $\mathcal{L}_{\Gamma, \mathbf{g}}$ as DG-algebras by direct computation of the Chekanov-Eliashberg DG-algebra of the Legendrian $L_{\Gamma, \mathbf{g}}$. In view of the surgery results from [5], [11], our result provides computations of various

wrapped Fukaya categories via an A_∞ quasi-isomorphism

$$\mathcal{B}_{\Gamma, \mathbf{g}} \cong \bigoplus_{v, w} CW^*(C_v, C_w)$$

where $CW^*(C_v, C_w)$ stands for the morphisms between the cocores C_v, C_w in the partially wrapped Fukaya of $(\#^{k+2g}(S^1 \times \mathbb{D}^3), \omega_{std})$ where all Legendrians L_v are used as stops (see [11, Conj. 3]). In this setting, it is important that the indeterminates t_v 's do not commute with the other generators. On the other hand, if one specialises to $t_v = e_v$ for $v = 1, \dots, s$, one recovers the fully wrapped Fukaya category of X_Γ where X_Γ is equipped with the exact symplectic form associated with its Weinstein structure. Setting the value of $\text{Log}(t_v) = \omega([S_v])$ to other constants, has the effect of turning on non-exact (bulk) deformations of the symplectic form, by giving area to the zero-sections S_v of the cotangent bundles involved in the plumbing X_Γ . One can interpret this as a computation of bulk deformed wrapped Fukaya categories.

Remark 3. As in [11], let

$$\mathcal{A}_{\Gamma, \mathbf{g}} = \bigoplus_{v, w} CF^*(S_v, S_w)$$

be the endomorphism algebra of the compact exact Lagrangians in X_Γ which form the Lagrangian skeleton of the plumbing X_Γ . After letting $t_v = e_v$, for $v = 1, \dots, s$, one has a canonical augmentation

$$\epsilon : \mathcal{B}_{\Gamma, \mathbf{g}} \rightarrow \mathbf{k}$$

corresponding to the module $\bigoplus CF^*(C_v, S_v)$ in the (fully) wrapped Fukaya category of $\mathcal{W}(X_\Gamma)$, given explicitly by

$$\epsilon(c_a) = \epsilon(c_a^*) = 0, \quad \epsilon(\alpha_{v,i}^\pm) = \epsilon(\beta_{v,i}^\pm) = e_v = \epsilon(z_a^\pm), \quad v = s(a)$$

By [11, Thm. 4], one can compute

$$\mathcal{A}_{\Gamma, \mathbf{g}} \simeq \text{Rhom}_{\mathcal{B}_{\Gamma, \mathbf{g}}}(\mathbf{k}, \mathbf{k})$$

This could be useful in studying Fukaya categories of symplectic 4-manifolds. Namely, if one has any configuration of transversely intersecting compact Lagrangians in a symplectic 4-manifold whose intersection pattern is encoded by the decorated graph (Γ, \mathbf{g}) , the corresponding A_∞ -algebra is a deformation of $\mathcal{A}_{\Gamma, \mathbf{g}}$.

We also note that the grading in $\mathcal{B}_{\Gamma, \mathbf{g}}$ is supported in non-positive degrees. In higher dimensional plumbings, Abouzaid and Smith used this property and an additional requirement on the zeroth cohomology of cocores [2, Prop. 4.2] that guarantees that the only modules supported in a single degree are given by the compact cores with local systems, to deduce that the compact cores of the plumbing generate the compact Fukaya category. In our case, this additional requirement does not hold. Nonetheless, the core Lagrangian surfaces $\{S_v\}$ are known to generate the compact Fukaya category $\mathcal{F}(X_\Gamma)$ in several situations, including when Γ is of Dynkin type with $\mathbf{g} = 0$ ([23, Lem. 4.15] and [24, Cor. 5.8]). We also remark that Keating [17] gave examples of Weinstein 4-manifolds which are *partial compactifications* of plumbings with $\mathbf{g} = 0$, where an exact Lagrangian torus (equipped with certain local systems) is not generated by the core spheres.

Remark 4. After we wrote our first paper [14] on computing wrapped Fukaya categories of X_Γ , which studied the case where Γ is a tree and $\mathbf{g} = 0$, an interesting paper by Bezrukavnikov and Kapranov [3] appeared, where finite dimensional $\Lambda_{\Gamma, \mathbf{g}}$ -modules are related to microlocal sheaves on the Lagrangian skeleton of X_Γ given by the union of transversely intersecting surfaces. In view of the correspondence between Fukaya categories of cotangent bundles and microlocal sheaves on the zero-section [19], Thm. (1) provides a confirmation of the more general expectation of an equivalence between Fukaya categories of Weinstein manifolds and microlocal sheaves on (singular) Lagrangian skeleta for the manifolds X_Γ . This more general equivalence is the subject of ongoing work by Ganatra, Pardon and Shende.

Let us consider the case Γ is a tree and $\mathbf{g} = 0$, and specialise to $t_v = e_v$. By definition of \mathcal{B}_Γ , the differential $\partial\tau_v$ consists of words of (even) length ≥ 2 . If we truncate it so that only words of length 2 appears, the resulting DG-algebra is known as the Ginzburg DG-algebra \mathcal{G}_Γ which was studied in [14]. It is known that if Γ is not of Dynkin type, \mathcal{G}_Γ has cohomology only in degree 0 (cf. Thm 8, Cor. 27 [14]) and $H^0(\mathcal{G}_\Gamma) = \Pi_\Gamma$ is the additive preprojective algebra. Now, by considering the word-length filtration, we get a spectral sequence from \mathcal{G}_Γ to \mathcal{B}_Γ . At least when $\text{char}\mathbb{K} = 0$, one can show that the length spectral sequence used in the above proof is degenerate. The reason for this was given in [14, pg. 7]. Namely, it is known that Π_Γ has no formal deformations as a Calabi-Yau algebra. However, the length spectral sequence does not converge strongly if Γ is non-Dynkin, hence one cannot conclude that \mathcal{B}_Γ and \mathcal{G}_Γ are quasi-isomorphic without having to complete both algebras. Indeed, for non-Dynkin Γ , the additive preprojective algebra $\Pi_\Gamma = H^0(\mathcal{G}_\Gamma)$ and the multiplicative preprojective algebra $\Lambda_\Gamma = H^0(\mathcal{B}_\Gamma)$ are not isomorphic. This can be seen by considering the representation varieties of Π_Γ and Λ_Γ - additive and multiplicative Nakajima quiver varieties. Thus, it is the (derived) multiplicative preprojective algebra \mathcal{B}_Γ that is central to the study of wrapped Fukaya categories of plumbings, not the additive preprojective algebra (or its derived version).

Example 5. When Γ is just a single vertex labelled with g , then X_Γ is just the cotangent bundle of a genus g surface Σ_g . In this case, we see that the multiplicative preprojective algebra is simply the group-ring $\mathbb{K}[\pi_1(\Sigma_g)]$, which agrees with the based loop space homology of the cotangent fibre. This agrees with the computation given in [12].

Example 6. When Γ has a single vertex (labelled with 0) and a single loop, the symplectic manifold X_Γ obtained by self-plumbing of T^*S^2 was studied in detail in [22]. In this case, it is easy to see that (cf. [8, Ex. 1.3]) the multiplicative preprojective algebra (with $t = 1$) Λ_Γ is quasi-isomorphic to the commutative algebra $\mathbb{K}[x, y, (1 + xy)^{-1}]$, which agrees with the homological mirror symmetry statement proven in [22] (cf. [6]). This variety has a holomorphic symplectic form given by $\Omega = dx \wedge dy / (1 + xy)$ which agrees with the standard holomorphic symplectic form on the two copies of $(\mathbb{C}^\times)^2$ given by the co-ordinates $(x, 1 + xy)$ (for $x \neq 0$) and $(y, 1 + xy)$ (for $y \neq 0$). One can consider the quantization of the functions compatible with the Poisson structure induced by Ω . We point out that if t is not specialized to 1 but carried along as a commuting complex parameter corresponding

to an area parameter on the immersed sphere, then we get the “quantized algebra” where x and y no longer commute but instead satisfy

$$1 + xy = t(1 + yx)$$

where $(1 + xy)$ is still invertible. Indeed, it is easy to see from this that we get the standard quantum torus algebras on each copy of $(\mathbb{C}^*)^2$. For example, we have

$$t(1 + xy)x = tx(1 + yx) = x(1 + xy)$$

We end the introduction by pointing out an intriguing connection to (multiplicative) Nakajima quiver varieties that we plan to return to in the future. Note that such a connection was also observed in [3] (see also [25]) via calculations using microlocal sheaves.

It is well known that the moduli spaces of representations of the multiplicative preprojective algebra Λ_Γ has the structure of algebraic symplectic varieties known as the multiplicative Nakajima quiver varieties (studied by Yamakawa [26], see also Boalch [4]). By our result, any Lagrangian $L \subset X_\Gamma$ gives a representation $\bigoplus_v \text{Hom}(C_v, L)$ of the multiplicative preprojective algebra $\bigoplus_{v,w} \text{Hom}(C_v, C_w)$. Thus, our result can be interpreted as the statement that “moduli space of objects” in the derived Fukaya category of X_Γ can be identified with the corresponding multiplicative Nakajima quiver varieties. These are finite type varieties once one fixes a dimension vector consisting of $d_v \in \mathbb{Z}_{\geq 0}$ for each vertex v of Γ and a stability condition encoded by $\theta_v \in \mathbb{Q}$ for each v , satisfying $\sum_v d_v \theta_v = 0$, and the values of $t_v \in \mathbb{C}^\times$. The choice of dimension vectors corresponds to fixing the rank $d_v = \dim_{\mathbb{K}} \text{Hom}(L, C_v)$, and the choice of stability condition should correspond to fixing a choice of Bridgeland stability on the derived Fukaya category of X_Γ . Finally, note that the symplectic structures on the Nakajima quiver varieties correspond to the fact that $\mathcal{B}_{\Gamma, \mathbf{g}}$ is 2-Calabi-Yau (cf. [21]).

In Section 2, after reviewing the construction of Chekanov-Eliashberg algebra associated to a Legendrian surgery presentation of a Weinstein 4-manifold following Ekholm-Ng [12], we provide a simplification that results in a finitely generated DG-algebra even in the presence of 1-handles. In Section 3, we draw a Legendrian surgery presentation of plumbings of T^*S^2 ’s according to an arbitrary graph and prove our main theorem for $\mathbf{g} = 0$. In Section 4, we extend the results of Section 3 to arbitrary \mathbf{g} .

Acknowledgments: T.E. is partially supported by the Technological and Research Council of Turkey through a BIDEB-2219 fellowship. Most of the work was carried out while T.E. was visiting Princeton University. Y.L. is supported in part by the Royal Society (URF) and the NSF grant DMS-1509141. We would like to thank Pavel Etingof, Lenhard Ng, and Ivan Smith for helpful correspondence. We also thank referees for their comments.

2. LEGENDRIAN DG-ALGEBRA ON A SUBCRITICAL WEINSTEIN 4-MANIFOLD

Recall that any Weinstein 4-manifold X has a Weinstein handlebody decomposition consisting of 0-, 1- and 2-handles. Assuming X is connected, one can find a handlebody decomposition with a unique 0-handle. Furthermore, Weinstein 1-handles can be attached

in a unique way, hence the more interesting part of the handlebody decomposition is the attachment of Weinstein 2-handles along Legendrian knots in $\#^k(S^1 \times S^2)$ equipped with the unique Stein-fillable contact structure, where k is the number of 1-handles. If $k = 0$, the 2-handles are attached along Legendrian knots in S^3 equipped with its standard contact structure.

Following Gompf [15], a surgery diagram for X is usually presented as a front diagram in a *standard form*. In order to give a combinatorial description of the Chekanov-Eliashberg's Legendrian DG-algebra ([7, 13]) associated with a Legendrian link $L \subset \#^k(S^1 \times S^2)$ given in Gompf's standard form, Ekholm and Ng [12] described a procedure called *resolution* (analogous to the resolution of a Legendrian in S^3 in [20]). We give an example in Figure 1 and refer the readers to the original references [15, 12] for precise definitions.




FIGURE 1. A Legendrian knot in Gompf normal form (top), Ekholm-Ng's resolution (middle), A simplifying Legendrian isotopy via Reidemeister III (bottom)

We next recall the description of the Legendrian DG-algebra, here denoted $CE^*(L)$, that Ekholm and Ng provide coming from a resolution diagram. All our complexes are cohomological, thus we reverse the gradings from [12].

Fix a field \mathbb{K} . Let $L = L_1 \cup L_2 \cup \dots \cup L_s \subset \#^k(S^1 \times S^2)$ be a link given in the resolved form. Let a_1, \dots, a_n denote the crossings of the resolution diagram. One refers to the set $\{a_1, \dots, a_n\}$ as the set of *external generators*. Note each a_i corresponds to a Reeb chord connecting L_v to L_w for some v and w , where L_v is the label of the under-strand and L_w is the label of the over-strand at the crossing a_i . Thus, we can form the \mathbf{k} -bimodule

$$\mathcal{E} := \bigoplus_{i=1}^n \mathbb{K}a_i$$

where $e_w a_i e_v = a_i$ precisely if a_i corresponds to a Reeb chord from L_v to L_w .

In the interior region of each 1-handle, we also have generators. These will be referred to as the *internal generators*. Following Ekholm-Ng, we describe these generators. Let k be the number of 1-handles. For each $1 \leq l \leq k$, let o_l denote the number of strands of L entering to l -th handle. We label the strands on the left of the diagram from top to bottom by $1, \dots, o_l$ and on the right of the diagram from bottom to top by $1, \dots, o_l$ (note the switch in the ordering). For each 1-handle, there are infinitely many internal generators. For $1 \leq l \leq k$, these are labelled as follows:

$$\begin{aligned} c_{ij;l}^0 & \text{ for } 1 \leq i < j \leq o_l, \\ c_{ij;l}^p & \text{ for } p > 0 \text{ and } 1 \leq i, j \leq o_l. \end{aligned}$$

The generator $c_{ij;l}^p$ for $p \geq 0$ connect the i^{th} strand to j^{th} strand going to the handle labelled l , thus corresponds to a Reeb chord from some L_v to another L_w . This defines a \mathbf{k} -bimodule structure by defining $e_w c_{ij;l}^p e_v = c_{ij;l}^p$. We write

$$\mathcal{I} = \mathcal{I}_1 \oplus \mathcal{I}_2 \oplus \dots \oplus \mathcal{I}_k$$

for the corresponding \mathbf{k} -bimodule generated by the internal generators.

Definition 7. We let $CE^*(L)$ to be the DG-algebra over \mathbf{k} generated by the external generators

$$a_1, a_2, \dots, a_n,$$

the internal generators

$$\begin{aligned} c_{ij;l}^0, & \ 1 \leq i < j \leq o_l, \ l = 1, \dots, k \\ c_{ij;l}^p, & \ p > 0, \ 1 \leq i, j \leq o_l, \ l = 1, \dots, k, \end{aligned}$$

and

$$t_1, t_1^{-1}, \dots, t_s, t_s^{-1}$$

with relations

$$\begin{aligned} t_v t_v^{-1} &= t_v^{-1} t_v = e_v, \text{ for all } v = 1, \dots, s, \\ t_v^{\pm} t_w^{\pm} &= 0, \text{ for } v \neq w \end{aligned}$$

and the \mathbf{k} -bimodule structure described above.

The differential on the external generators is given by

$$\partial(a_i) = \sum_{r \geq 0} \sum_{b_1, \dots, b_r} \sum_{\Delta \in \Delta(a_i; b_1, \dots, b_r)} \pm \{t_1^{-n_1(\Delta)}, \dots, t_s^{-n_s(\Delta)}, b_r, \dots, b_1\}$$

where b_q are either external generators or of the form $c_{ij;l}^0$ and n_v record the number of times the boundary of the holomorphic disk Δ passed through $*_v$, and the notation

$$\{t_1^{n_1(\Delta)}, \dots, t_s^{n_s(\Delta)}, b_r, \dots, b_1\}$$

stands for the time-ordered product where reading from right-to-left corresponds to the order of terms appearing in the boundary of Δ when it is traversed counter-clockwise. In

particular, note that t_v and b_q maybe interlaced, as we do not necessarily insist that these commute. See [12, Sec. 2.4C] for the determination of the sign.¹

The differential on the internal generators (see Def. (8)) satisfies $\partial(\mathcal{I}_l) \subset \mathcal{I}_l$ for all $l = 1, \dots, k$ and only depends on the number of strands passing through the 1-handle. Thus, the internal DG-algebra corresponding to a 1-handle with n strands is denoted by \mathcal{J}_n .

The differential of a generators of type t_v is trivial.

It is possible to put a \mathbb{Z} -grading on CE^* (by making some extra choices). We refer to [12, Sec. 2.4B] for a detailed discussion of gradings on CE^* . Though, we use the negative of the grading given in [12] so that the differential increases this grading by 1. Note that the grading $|t_v| = 2r(L_v)$ where $r(L_v)$ is the rotation number.

We shall next discuss the internal algebra \mathcal{J}_n of Ekholm-Ng in more detail and show that it can be simplified to only include the free generators of the form $c_{ij;l}^0$ and $c_{ij;l}^1$. Hence, in particular, it is finitely generated up to quasi-isomorphism.

2.1. A simplification of Ekholm-Ng's internal DG-algebra. In Ekholm-Ng [12], the internal algebra \mathcal{J}_n was constructed geometrically from the study of Reeb chords internal to a 1-handle. It has the undesirable feature of being infinitely generated. Here, we provide smaller models for the quasi-isomorphism type of \mathcal{J}_n . In particular, we compute the cohomology of \mathcal{J}_n for all n , and show that \mathcal{J}_n is quasi-isomorphic to a finitely generated subalgebra of itself.

Note that throughout this paper the base ring is taken to be the semisimple ring \mathbf{k} where the number of idempotents is given by the number of 2-handles in the surgery diagram corresponding to the components of the Legendrian link.

In order to achieve full generality, we shall first define \mathcal{J}_n as a DG-algebra over a different ring

$$\mathbf{k}_n = \bigoplus_{i=1}^n \mathbb{K}e_i$$

As usual, n denotes the number of strands entering the 1-handle, irrespective of the 2-handles that these strands belong to. On the other hand, the internal DG-subalgebra of $CE^*(L)$ for a particular L is obtained from \mathcal{J}_n by a base change $\mathbf{k} \rightarrow \mathbf{k}_n$ given by the configuration of 2-handles entering in the relevant 1-handle. More precisely, at each 1-handle, the ring map from $\mathbf{k} \rightarrow \mathbf{k}_n$ is given by sending the idempotent e_v associated to a 2-handle L_v to the sum of idempotents $e_{i_1} + \dots + e_{i_v}$ in \mathbf{k} where i_1, \dots, i_v are the strands that pass through the 1-handle which belong to 2-handle L_v . For example, if a component of L_v enters to the same 1-handle more than once, then a chord that goes between the two strands of L_v can be composed with itself (arbitrarily many times).

We note that in [12], \mathbf{k} is always taken to be the ground field \mathbb{K} , regardless of the number of components of L entering in the 1-handle. This is the opposite extreme to taking $\mathbf{k} = \mathbf{k}_n$

¹As in [14], [11], we follow the convention that multiplications are read from *right-to-left*. This is not the same convention used in [12]. Throughout, our formulae are adjusted accordingly by taking the opposite DG-category (see [14, pg. 8]).

in that all the chords between any strands are allowed to be composable. However, from the perspective of Legendrian surgery, the correct choice if L is not connected is to work over a ring \mathbf{k} with idempotents in bijection with the connected components of L .

We give a formal definition as follows:

Definition 8. We define \mathcal{J}_n to be a DG-category over \mathbb{K} with n -objects : X_1, \dots, X_n . Morphism spaces are spanned (as vector spaces over \mathbb{K}) by

- the identity morphisms $e_i \in \text{hom}(X_i, X_i)$,
- the “elementary” morphisms in $\text{hom}(X_i, X_j)$

$$c_{ij}^0 : 1 \leq i < j \leq n \quad \text{and} \quad c_{ij}^p : 1 \leq p, 1 \leq i, j \leq n ,$$

and all possible compositions of the elementary morphisms, e.g. $c_{kl}^r c_{jk}^q c_{ij}^p \in \text{hom}(X_i, X_l)$. The (cohomological) grading is defined by first fixing n integers (m_1, m_2, \dots, m_n) and then defining:

$$|c_{ij}^p| = 1 - 2p + m_j - m_i.$$

The differential ∂ is determined by

$$\begin{aligned} \partial c_{ij}^0 &= \sum_{i < r < j} (-1)^{m_r + m_j} c_{rj}^0 c_{ir}^0, \quad i < j \\ \partial c_{ij}^1 &= \delta_{ij} + \sum_{r > i} (-1)^{m_r + m_j} c_{rj}^1 c_{ir}^0 + \sum_{r < j} (-1)^{m_r + m_j} c_{rj}^0 c_{ir}^1 \\ \partial c_{ij}^p &= \sum_{0 \leq q \leq p} \sum_r (-1)^{m_r + m_j} c_{rj}^{p-q} c_{ir}^q, \quad p > 1 \end{aligned}$$

(c_{ij}^0 is set to be 0 for $i \geq j$, $\delta_{ij} = e_i = e_j$ if $i = j$, and 0 otherwise) and the graded Leibniz rule:

$$\partial(a_2 a_1) = \partial(a_2) a_1 + (-1)^{|a_2|} a_2 \partial(a_1).$$

We define \mathcal{A}_n to be the DG-subcategory of \mathcal{J}_n with the same objects X_1, \dots, X_n , but a restricted set of elementary morphisms

$$c_{ij}^0 : 1 \leq i < j \leq n \quad \text{and} \quad c_{ij}^1 : 1 \leq i, j \leq n.$$

As is customary, we use the same notation for a DG-category \mathcal{C} and its total (endomorphism) algebra $\oplus_{X, Y \in \text{Ob } \mathcal{C}} \text{hom}(X, Y)$ which is a DG-algebra over the corresponding semisimple ring. For example, \mathcal{J}_n and \mathcal{A}_n denote DG-algebras over \mathbf{k}_n . On the other hand, given a unital ring map $\mathbf{k} \rightarrow \mathbf{k}_n$, $\mathcal{J}_n^{\mathbf{k}}$ and $\mathcal{A}_n^{\mathbf{k}}$ denote the corresponding DG-algebras over \mathbf{k} . The differential on $\mathcal{J}_n^{\mathbf{k}}$ (and similarly $\mathcal{A}_n^{\mathbf{k}}$) given by the same formulae as for \mathcal{J}_n (and similarly for \mathcal{A}_n) except that now some of the objects are identified, making more compositions possible, and the DG-algebra is generated by the same set of elementary morphisms over \mathbf{k} rather than \mathbf{k}_n . For example, in the case $\mathbf{k} = \mathbb{K}$ (as considered in [12]) arbitrary words in c_{ij}^p are allowed since all of them belong to the morphism space of the single object of the DG-category. We warn the reader that $\mathcal{J}_n^{\mathbb{K}}$ is quite different than considering \mathcal{J}_n and viewing it as a DG-algebra over \mathbb{K} by forgetting its \mathbf{k}_n -algebra structure.

A helpful way of thinking about these categories is to draw a circle with n points ordered in counter-clockwise direction corresponding to the objects X_i as in Figure 2. The set of generators c_{ij}^p is in bijection with homotopy classes of paths following a counter-clockwise flow that connects i to j . The exponent p counts the number of times such paths pass from an additional marked point $*$ between n and 1. Multiplication in the category can be thought of as broken paths, and the differential corresponds to breaking the path into two paths of shorter length, with the exception that in the case of ∂c_{ii}^1 , there is an additional idempotent term e_i that appears in the differential.




FIGURE 2. Visualisation of generators c_{ij}^p .

Next, let us introduce the following A_∞ -algebra, whose definition is attributed to Kontsevich in [24, Rmk. 3.11].

Definition 9. We define \mathcal{K}_n to be the strictly unital A_∞ -category with n -objects Z_1, \dots, Z_n and morphisms given by, for $i \in \mathbb{Z}/n$,

$$\begin{aligned} \text{hom}(Z_i, Z_i) &= \mathbb{K}e_i, \\ \text{hom}(Z_i, Z_{i+1}) &= \mathbb{K}x_i, \end{aligned}$$

where e_i are idempotents. We again fix n integers (m_1, m_2, \dots, m_n) and define gradings

$$|x_i| = 1 + m_{i+1} - m_i \text{ for } i = 1, \dots, n-1, \text{ and } |x_n| = -1 + m_1 - m_n$$

The A_∞ -operations involving idempotents are determined by strict unitality. The only other products are given by:

$$\mathbf{m}_n(x_{i-1}, \dots, x_1, x_0, x_{n-1}, \dots, x_{i+1}, x_i) = e_i, \text{ for } i \in \mathbb{Z}/n$$

The fact that A_∞ -relations are satisfied is checked easily by the following equation that holds for all $i \in \mathbb{Z}/n$:

$$\begin{aligned} &\mathbf{m}_2(x_i, \mathbf{m}_n(x_{i-1}, \dots, x_1, x_0, x_{n-1}, \dots, x_{i+1}, x_i)) \\ &+ (-1)^{|x_i|-1} \mathbf{m}_2(\mathbf{m}_n(x_i, \dots, x_1, x_0, x_{n-1}, \dots, x_{i+1}), x_i) = x_i - x_i = 0 \end{aligned}$$

Notice that we used $\mathbf{m}_2(e_{i+1}, x_i) = (-1)^{|x_i|} x_i$ (cf. [24, Eq. (1.3)]).

It can be shown that for $n \geq 3$, \mathcal{K}_n is not formal as an A_∞ -algebra, but we have

$$\mathcal{K}_2 \simeq \mathbf{k}_2 \langle x_1, x_2 \rangle / \langle x_2 x_1 = e_1, x_1 x_2 = e_2 \rangle \quad |x_1| = 1 + m_2 - m_1, |x_2| = -1 + m_1 - m_2$$

Recall that cobar and bar constructions extend to the setting of A_∞ -algebras. We denote these by Ω and B , respectively. (See [11, Sec. 2] for a review that is compatible with our sign conventions.)

Lemma 10. *\mathcal{I}_n is isomorphic to $\Omega B\mathcal{K}_n$. In particular, there exists a (strictly unital) A_∞ -functor*

$$\mathfrak{f} : \mathcal{K}_n \rightarrow \mathcal{I}_n$$

which is a quasi-isomorphism.

Proof. Recall that for any coaugmented conilpotent DG-coalgebra \mathcal{C} and an augmented DG-algebra \mathcal{A} we have an adjunction

$$\mathrm{hom}_{DG}(\Omega\mathcal{C}, \mathcal{A}) \simeq \mathrm{hom}_{coDG}(\mathcal{C}, B\mathcal{A})$$

(see [11, Sec. 2.2.2] for a recent exposition of this well-known adjunction.) In particular, for any A_∞ -algebra \mathcal{A} , applying the adjunction

$$\mathrm{hom}_{DG}(\Omega B\mathcal{A}, \Omega B\mathcal{A}) \simeq \mathrm{hom}_{coDG}(B\mathcal{A}, B\Omega B\mathcal{A})$$

and tracing the image of the identity map $\Omega B\mathcal{A} \rightarrow \Omega B\mathcal{A}$, we get a quasi-isomorphism of DG-coalgebras $B\mathcal{A} \rightarrow B\Omega B\mathcal{A}$, or equivalently an A_∞ -algebra quasi-isomorphism

$$\mathfrak{f} : \mathcal{A} \rightarrow \Omega B\mathcal{A}$$

for any A_∞ -algebra \mathcal{A} . This is sometimes referred to as the standard resolution of A (see for ex. [10, Sec. 16.7.3].

Chasing through the adjunction (see [11, Sec. 2.2.2]), one sees that the A_∞ functor $\mathfrak{f} = (\mathfrak{f}^i)_{i \geq 1}$ has an explicit description: A composable word consisting of i letters in \mathcal{A} is sent by \mathfrak{f}^i to the corresponding element of $B\mathcal{A}$ (whose elements by definition consist of words in \mathcal{A}).

Here, we apply this construction to $\mathcal{A} = \mathcal{K}_n$. Indeed, the A_∞ -functor \mathfrak{f} is the Yoneda embedding defined by sending $\mathfrak{f}(Z_i) = X_i$ at the level of objects, and on morphisms, we set for $k \geq 1$:

$$\mathfrak{f}(x_{i+k-1}, x_{i+k-2}, \dots, x_i) = (-1)^{|c_{ij}^p|} c_{ij}^p$$

where j and p are determined uniquely by the equation $i + k = j + np$, $1 \leq j \leq n$. Here, the indices i appearing on generators x_i are considered in \mathbb{Z}/n . The A_∞ -functor equation takes the form (using the rules [14, Eq. 4]):

$$\sum_{0 \leq q \leq p} \sum_r (-1)^{|c_{ir}^q|} \mathfrak{f}(x_{j+np-1}, \dots, x_{r+nq}) \mathfrak{f}(x_{r+nq-1}, \dots, x_i) + \partial c_{ij}^p = \delta_{ij} \delta_{1p}$$

which coincides precisely with the definition of the differential on \mathcal{I}_n .

Note that in [11, Sec. 2], the signs for the cobar-bar constructions are written out explicitly. It is an easy exercise to check that the signs are compatible with the signs that appear in Definition 8 of the internal algebra.

Finally, the fact that \mathfrak{f} is quasi-isomorphism follows from the bar-cobar adjunction as explained above. \square

Corollary 11. $H^*(\mathcal{J}_n)$ is generated over \mathbf{k}_n by the elements

$$c_{12}^0, c_{23}^0, \dots, c_{(n-1)n}^0, c_{n1}^1.$$

This computation can be used to show that the DG-subalgebra \mathcal{A}_n of \mathcal{J}_n generated by only the elements

$$c_{ij}^0 : 1 \leq i < j \leq n \text{ and } c_{ij}^1 : 1 \leq i, j \leq n$$

is quasi-isomorphic to \mathcal{J}_n . Here we provide a general proof over any base ring \mathbf{k} .

Theorem 12. For every base ring \mathbf{k} , let $\mathbf{i} : \mathcal{A}_n^{\mathbf{k}} \rightarrow \mathcal{J}_n^{\mathbf{k}}$ be the inclusion map. There exists a retraction $\mathbf{r} : \mathcal{J}_n^{\mathbf{k}} \rightarrow \mathcal{A}_n^{\mathbf{k}}$ given by

$$\begin{aligned} \mathbf{r}|_{\mathcal{A}_n^{\mathbf{k}}} &= \text{Id}_{\mathcal{A}_n^{\mathbf{k}}} \\ \mathbf{r}(c_{ij}^p) &= C_{p-1} \sum_{k_1, \dots, k_{2p-2}} (-1)^{\dagger} c_{k_{2p-2}j}^1 \cdots c_{k_1 k_2}^1 c_{ik_1}^1 \text{ for } p > 1, \end{aligned}$$

where $C_p = \frac{1}{p+1} \binom{2p}{p}$ is the p^{th} Catalan number and $\dagger = m_{k_1} + \dots + m_{k_{2p-2}}$, that induces a quasi-isomorphism. The latter follows from the existence of a chain homotopy \mathfrak{H} between $\mathbf{i} \circ \mathbf{r} : \mathcal{J}_n^{\mathbf{k}} \rightarrow \mathcal{J}_n^{\mathbf{k}}$ and the identity map defined by

$$\mathfrak{H}|_{\mathcal{A}_n^{\mathbf{k}}} = 0 \tag{1}$$

$$\mathfrak{H}(c_{ij}^p) = \sum_{q=2}^p C_{p-q+1} \sum_{k_1, \dots, k_{2p-2q+1}} (-1)^{\ddagger} c_{k_{2p-2q+1}j}^q c_{k_{2p-2q} k_{2p-2q+1}}^1 \cdots c_{k_1 k_2}^1 c_{ik_1}^1 \text{ for } p > 1 \tag{2}$$

$$\mathfrak{H}(a_2 a_1) = \mathfrak{H}(a_2) \mathbf{r}(a_1) + (-1)^{|a_2|} a_2 \mathfrak{H}(a_1) \tag{3}$$

where $\ddagger = m_{k_1} + m_{k_2} + \dots + m_{k_{2p-2q+1}} + m_j$.

Proof. First, we check that \mathbf{r} is a DG-algebra map by verifying that \mathbf{r} commutes with the differential on the generators.

For $p > 1$,

$$\begin{aligned} \mathbf{r} \partial(c_{ij}^p) &= \sum_{k_1, k_2, \dots, k_{2p-1}} (-1)^{\dagger + m_j + m_{k_{2p-1}}} C_{p-1} \left(c_{k_{2p-1}j}^0 c_{k_{2p-2} k_{2p-1}}^1 \cdots c_{ik_1}^1 + c_{k_{2p-1}j}^1 \cdots c_{k_1 k_2}^1 c_{ik_1}^0 \right) \\ &+ \sum_{k_1, k_2, \dots, k_{2p-3}} (-1)^{\dagger + m_j - m_{k_{2p-2}}} (C_{p-2} + C_2 C_{p-3} + \cdots + C_{p-3} C_2 + C_{p-2}) c_{k_{2p-3}j}^1 \cdots c_{ik_1}^1 \end{aligned}$$

whereas

$$\begin{aligned} \partial \mathbf{r}(c_{ij}^p) &= C_{p-1} \left(\sum_{k_1, k_2, \dots, k_{2p-1}} (-1)^{\dagger + m_j + m_{k_{2p-1}}} \left(c_{k_{2p-1}j}^0 c_{k_{2p-2} k_{2p-1}}^1 \cdots c_{ik_1}^1 + c_{k_{2p-1}j}^1 \cdots c_{k_1 k_2}^1 c_{ik_1}^0 \right) \right. \\ &\quad \left. + \sum_{k_1, k_2, \dots, k_{2p-3}} (-1)^{\dagger + m_j - m_{k_{2p-2}}} c_{k_{2p-3}j}^1 \cdots c_{ik_1}^1 \right) \end{aligned}$$

after cancellations. The fact that Catalan numbers satisfy the recurrence relation

$$C_{p-1} = \sum_{q=0}^{p-2} C_q C_{p-q-2}$$

shows that \mathfrak{r} is a DG-algebra homomorphism.

Next, we prove that \mathfrak{H} is a chain homotopy between $\mathfrak{i} \circ \mathfrak{r}$ and the identity map. Observe that, by (3), it suffices to check the condition $\partial \mathfrak{H} + \mathfrak{H} \partial = \mathfrak{i} \circ \mathfrak{r} - \text{Id}$ on the generating set $\{c_{ij}^p\}$ which is clearly satisfied for $p = 0, 1$. From now on let $p \geq 2$.

In order to compute $(\partial \mathfrak{H} + \mathfrak{H} \partial)(c_{ij}^p)$, first we have

$$\mathfrak{H} \partial(c_{ij}^p) = \sum_{q=0}^p \sum_k (-1)^{m_k + m_j} \mathfrak{H}(c_{kj}^q c_{ik}^{p-q})$$

and, by (3), this is a sum of two expressions, first of which is

$$\sum_{q=2}^{p-1} \sum_{r=2}^q C_{q-r+1} C_{p-q-1} \sum_{k_s} (-1)^\epsilon c_{k_{2p-2r}j}^r c_{k_{2p-2r-1}k_{2p-2r}}^1 \cdots c_{ik_1}^1 \quad (4)$$

$$+ \sum_{r=2}^p C_{p-r+1} \sum_{k_s} (-1)^\epsilon c_{k_{2p-2r+2}j}^r c_{k_{2p-2r-1}k_{2p-2r}}^1 \cdots c_{k_1 k_2}^1 c_{ik_1}^0 \quad (5)$$

and the second is

$$\sum_{q=0}^p \sum_{r=2}^{p-q} C_{p-q-r+1} \sum_{k_s} (-1)^{1+\epsilon} c_{k_{2p-2q-2r+2}j}^q c_{k_{2p-2q-2r+1}k_{2p-2q-2r+2}}^r c_{k_{2p-2q-2r}k_{2p-2q-2r+1}}^1 \cdots c_{ik_1}^1 \quad (6)$$

$$= \sum_{q'=2}^p \sum_{r=2}^{q'} C_{p-q'+1} \sum_{k_s} (-1)^{1+\epsilon} c_{k_{2p-2q'+2}j}^{q'-r} c_{k_{2p-2q'+1}k_{2p-2q'+2}}^r c_{k_{2p-2q'}k_{2p-2q'+1}}^1 \cdots c_{ik_1}^1 \quad (7)$$

where $\epsilon = \sum_s m_{k_s}$. Note that the expression (7) (which will be cancelled by a term below) is obtained from expression (6) by a change of variable $q' = q + r$.

We also have

$$\partial \mathfrak{H}(c_{ij}^p) = \sum_{q'=2}^p C_{p-q'+1} \sum_l (-1)^\dagger \partial \left(c_{l_{2p-2q'+1}j}^{q'} c_{l_{2p-2q'}l_{2p-2q'+1}}^1 \cdots c_{l_1 l_2}^1 c_{il_1}^1 \right)$$

and, by the graded Leibniz rule, this is a sum of two expressions, first of which is based on $\left(\partial c_{l_{2p-2q'+1}j}^{q'}\right) \cdot c_{l_{2p-2q'}l_{2p-2q'+1}}^1 \cdots c_{l_1l_2}^1 c_{il_1}^1$

$$\sum_{q'=2}^p C_{p-q'+1} \sum_{r=1}^{q'} \sum_{k_s} (-1)^\epsilon c_{k_{2p-2q'}+2j}^{q'-r} c_{k_{2p-2q'}+1}^r c_{k_{2p-2q'}+2}^1 c_{k_{2p-2q'}k_{2p-2q'+1}}^1 \cdots c_{ik_1}^1 \quad (8)$$

$$+ \sum_{q'=2}^p C_{p-q'+1} \sum_{k_s} (-1)^\epsilon c_{k_{2p-2q'}+2j}^{q'} c_{k_{2p-2q'}+1}^0 c_{k_{2p-2q'}+2}^1 c_{k_{2p-2q'}k_{2p-2q'+1}}^1 \cdots c_{ik_1}^1 \quad (9)$$

Note that $r \geq 2$ part of (8) cancels (7) above and (9) is cancelled by a sum in the second part of $\mathfrak{H}\partial(c_{ij}^p)$ whose remaining contributions are as follows

$$\sum_{q'=2}^p C_{p-q'+1} \sum_{k_s} (-1)^{1+\epsilon} c_{k_{2p-2q'}j}^{q'} c_{k_{2p-2q'}-1}^1 c_{k_{2p-2q'}}^1 \cdots c_{ik_1}^1 \quad (10)$$

$$+ \sum_{q'=2}^p C_{p-q'+1} \sum_{k_s} (-1)^{1+\epsilon} c_{k_{2p-2q'}+2j}^r c_{k_{2p-2q'}-1}^1 c_{k_{2p-2q'}}^1 \cdots c_{k_1k_2}^1 c_{ik_1}^0 \quad (11)$$

where (10) is a result of the Kronecker delta component in the differential of c_{kl}^1 . Observe that (11) is cancelled by (5) and therefore $(\partial\mathfrak{H} + \mathfrak{H}\partial)(c_{ij}^p)$ is equal to the sum of the expressions (4), (10), and what remains from (8) after cancellation by (7), namely

$$\sum_{q'=2}^p C_{p-q'+1} \sum_{k_s} (-1)^\epsilon c_{k_{2p-2q'}+2j}^{q'-1} c_{k_{2p-2q'}+1}^1 c_{k_{2p-2q'}+2}^1 \cdots c_{ik_1}^1$$

whose $q' = 2$ term is precisely $\mathfrak{r}(c_{ij}^p)$. Also note that $q' = p$ term of (10) is exactly c_{ij}^p . Therefore $(\partial\mathfrak{H} + \mathfrak{H}\partial)(c_{ij}^p)$ is equal to $(\mathfrak{i} \circ \mathfrak{r} - \text{Id})(c_{ij}^p)$ plus the following (which we obtain by switching the order of summation between q and r variables in (4) and renaming the variables q' in (8) and (10) as $r+1$ and r , respectively):

$$\sum_{r=2}^{p-1} \left(\left(\sum_{q=r}^{p-1} C_{q-r+1} C_{p-q-1} \right) - C_{p-r+1} + C_{p-r} \right) \sum_{k_s} (-1)^\epsilon c_{k_{2p-2r}j}^r c_{k_{2p-2r}-1}^1 c_{k_{2p-2r}}^1 \cdots c_{ik_1}^1$$

But the above expression vanishes since

$$C_{p-r+1} = \left(\sum_{q=r}^{p-1} C_{q-r+1} C_{p-q-1} \right) + C_{p-r}$$

by the recurrence relation satisfied by the Catalan numbers. \square

Remark 13. For the readers who prefer a less computational approach, we outline another proof that uses more abstract machinery but is less explicit.

The category \mathcal{K}_{n+1} appears as the partially wrapped Fukaya category of the \mathbb{D}^2 with $(n+1)$ marked points, see [16, Sec. 3.3]. In Figure 3, one can find the Lagrangians in this




FIGURE 3. \mathcal{K}_{n+1} represented as the partially wrapped category of \mathbb{D}^2 with $(n+1)$ marked points.

category that correspond to the objects Z_i for $i = 1, \dots, n+1$. As noted in [16, Sec. 3.3], one can easily check that the object $Z_{n+1}[-|x_{n+1}|]$ is isomorphic to the twisted complex:

$$Z_1 \xrightarrow{x_1} Z_2[|x_1| - 1] \xrightarrow{x_2} \dots \xrightarrow{x_{n-1}} Z_n[|x_1| + |x_2| + \dots + |x_{n-1}| + 1 - n]$$

where the isomorphism maps are given by x_n and x_{n+1} .

Thus, if we take the full subcategory of \mathcal{K}_{n+1} consisting of objects Z_1, \dots, Z_n and localise it along the above twisted complex representing Z_{n+1} , we should obtain \mathcal{K}_n (as shown in [16, Prop. 3.5], localising along the object Z_{n+1} corresponds to the removal of the marked point near which it is supported).

Now, it is an easy exercise in localisation of DG-categories ([10], cf. [16, Sec 3.5]) to show that localisation of Z_1, \dots, Z_n along the above twisted complex can be identified directly with \mathcal{A}_n (cf. proof of Prop. 14 below).

We note that cohomology of \mathcal{I}_n and $\mathcal{I}_n^{\mathbf{k}}$ can be very different. In what follows we will use the following result that follows from Theorem 12 and the Proposition 14 given below :

$$\mathcal{I}_2^{\mathbb{K}} \simeq \mathbb{K}[z, z^{-1}] \quad (12)$$

In particular, note that this is infinite-dimensional over \mathbb{K} . On the other hand, as we proved in Corollary 11, cohomology of \mathcal{I}_2 is generated by two elements over \mathbf{k}_2 .

Proposition 14. $\mathcal{A}_2^{\mathbb{K}}$ is quasi-isomorphic to $\mathbb{K}[z, z^{-1}]$ considered with trivial differential.

Proof. To simplify the notation, let us denote $c_{12}^0, c_{21}^1, c_{11}^1, c_{22}^1$, and c_{12}^1 by s, t, k, l , and u , respectively. Thus, the algebra $\mathcal{A}_2^{\mathbb{K}}$ is the semifree DG-algebra, generated over \mathbb{K} freely by s, t, k, l, u with the differential determined by

$$\begin{aligned} dk &= 1 - ts \\ dl &= 1 - st \\ du &= ls - sk \end{aligned}$$

where the gradings are determined as follows $|s| = m, |t| = -m, |k| = |l| = -1, |u| = m - 2$ for an arbitrary fixed integer m . We would like to show that this DG-algebra is quasi-isomorphic to the algebra $\mathbb{K}[z, z^{-1}]$ with $|z| = m$ considered as a DG-algebra with trivial

differential. We did not manage to construct a quasi-isomorphism explicitly even though, surely, such explicit construction is feasible. Instead, we will exploit the fact that the algebra $\mathcal{A}_2^{\mathbb{K}}$ is given by Drinfeld's construction of DG-quotient (as in [10]). Namely, let us consider the derived category $\mathcal{A} = D^b(\mathbb{K}[z])$. Let L denote the object corresponding to $\mathbb{K}[z]$, and $S = \text{Cone}(L[-m] \xrightarrow{z} L)$, and \mathcal{A}_0 be the thick triangulated subcategory of \mathcal{A} generated by S . Equivalently, \mathcal{A}_0 corresponds to modules of $\mathbb{K}[z]$ with support at 0. By the main result of [18, Thm. 3.2] Verdier quotient $\mathcal{A}/\mathcal{A}_0$ is identified with $D^b(\mathbb{K}[z, z^{-1}])$ and the endomorphism algebra of L in this quotient is $\text{End}_{\mathcal{A}/\mathcal{A}_0}(L) = \mathbb{K}[z, z^{-1}]$. On the other hand, by viewing $\mathbb{K}[z]$ as DG-algebra with trivial differential, we can consider the DG-enhancement of \mathcal{A} , call it \mathcal{A}^{pre} . We can then apply Drinfeld's construction given in [10, Sec. 3.1] and take the DG quotient of \mathcal{A}^{pre} by the full DG-subcategory \mathcal{A}_0^{pre} consisting of S (see [10, Ex. 3.7.1] for a similar looking example corresponding to $\mathcal{A}_2^{\mathbb{K}_2}$). The endomorphism algebra of L in $\mathcal{A}^{pre}/\mathcal{A}_0^{pre}$ is generated by s, t, k, l, u where these generators can be identified with the morphisms of twisted complexes given in Figure 4. Moreover, the differential of these generators are easily computed to give an identification of $\text{end}_{\mathcal{A}^{pre}/\mathcal{A}_0^{pre}}(L)$ with $\mathcal{A}_2^{\mathbb{K}}$. Now, by the general theorem of Drinfeld [10, Thm. 3.4], the homotopy categories of these two different quotients are isomorphic. In particular,

$$H^*(\text{end}_{\mathcal{A}^{pre}/\mathcal{A}_0^{pre}}(L)) \simeq \text{End}_{\mathcal{A}/\mathcal{A}_0}(L).$$

It follows that

$$H^*\mathcal{A}_2^{\mathbb{K}} = \mathbb{K}[z, z^{-1}]$$

Finally, it is well known that $\mathbb{K}[z, z^{-1}]$ is intrinsically formal for any value of m . Since one can easily compute $HH^*(\mathbb{K}[z, z^{-1}]) = \mathbb{K}[z, z^{-1}, \partial_z]$ and check that $HH_{<0}^2(\mathbb{K}[z, z^{-1}]) = 0$. Therefore, it follows that $\mathcal{A}_2^{\mathbb{K}}$ is quasi-isomorphic to $\mathbb{K}[z, z^{-1}]$. Note that for $m = 0$, we can argue more directly by noting that the cohomology is supported in degree 0, hence all the higher products vanish for degree reasons.




FIGURE 4. Primitive generators in the Drinfeld quotient, identified with s, t, k, l, u from left-to-right

□

Remark 15. Another proof of the Proposition 14 can be given by using partially wrapped Fukaya categories. Namely, consider a cylinder T^*S^1 with a single stop along one of its boundary components. The cotangent fiber L generates this category and one has $\text{End}(L) = \mathbb{K}[z]$. On the other hand, the object supported near the stop is quasi-isomorphic to $\text{Cone}(L \xrightarrow{z} L)$, localizing with respect to that object corresponds to removing the stop (by [16, Prop. 3.5]). Thus, we can identify the endomorphism algebra of L in the fully wrapped Fukaya category of T^*S^1 as this localisation which (as we showed in the proof of Proposition 14) coincides with $\mathcal{A}_2^{\mathbb{K}}$. Now, one can directly compute the endomorphism algebra of L in the fully wrapped Fukaya category of T^*S^1 , which gives $\mathbb{K}[z, z^{-1}]$.

3. PLUMBINGS OF COTANGENT BUNDLES OF 2-SPHERES

Throughout this section, we assume that Γ is a finite graph with $\mathbf{g} = 0$. We first construct a Legendrian link L_Γ in $\#^k(S^1 \times S^2, \xi_{std})$ associated to Γ . An important feature of this construction is that the symplectic 4-manifold obtained by adding symplectic 2-handles along L_Γ to $\#^k(S^1 \times \mathbb{D}^3, \omega_{std})$ is homotopic, as a Weinstein manifold, to the plumbing of copies of T^*S^2 with respect to the graph Γ .


After the construction, we compute the Chekanov-Eliashberg DG-algebra $\mathcal{B}_\Gamma = CE^*(L_\Gamma)$ of L_Γ and prove that it is quasi-isomorphic to the derived multiplicative preprojective algebra associated to Γ .

3.1. A Legendrian surgery presentation of plumbings. Choose a spanning tree T of Γ and embed it in the upper half plane $\mathbb{R} \times \mathbb{R}_{\geq 0}$ such that all the vertices lie on $\mathbb{R} \times \{0\}$. This can be easily done recursively (in multiple ways). We give an illustrating picture in Fig. (5) and leave the elaboration of the recursive argument to the reader.

Corresponding to the spanning tree T , we will have 2-handles in the Legendrian surgery picture of X_Γ . The complement of T will correspond to 1-handles.

Before we put the graph Γ in a standard form, we choose an orientation and order on the edges of Γ . Note that, as is the case for the multiplicative preprojective algebra Λ_Γ , \mathcal{B}_Γ is also independent of these choices. Then for each arrow $h \in \Gamma \setminus T$ we introduce a pair of arcs in the lower half plane $\mathbb{R} \times \mathbb{R}_{< 0}$ as in Fig. (5). More precisely, we fix m, M as real numbers smaller and larger than the x -coordinates of v_1 and v_s , respectively, and if h is an arrow from v to w which is the i^{th} arrow in $\Gamma \setminus T$, then there will be a curve from v to the point $(m, -i)$ with nonnegative slope throughout and another curve from w to the point $(M, -i)$ with nonpositive slope, chosen so that there are no triple intersections. We assume that the arrows in T are oriented from right to left and also assume that in the ordering of all the edges of Γ , the edges in T are ahead of the edges in $\Gamma \setminus T$.

Once Γ is in a standard form, the endpoints of each pair of arcs corresponding to the same arrow in $\Gamma \setminus T$ is replaced by the feet, i.e. the attaching sphere $S^0 \times D^3$, of a 1-handle and each vertex v is replaced by the standard Lagrangian projection of a Legendrian unknot U_v with $tb = -1$. For each arrow a from v to w , we introduce an arc on the left half of U_v and a meridian m_a of the right half of U_w , i.e. the boundary of a small disk intersecting it transversely once, (so that all meridians and arcs on an unknot are ordered according to the chosen order of the edges), replace a by a band from the arc on U_v to an arc on m_a , and

FIGURE 5. The graph Γ in a standard form with respect to a chosen orientation

then connect $\text{sum } U_v$ and m_a using this band. None of these bands contain a twist other than the one near the left foot of the corresponding 1-handle (dictated by the resolution operation [12, Def. 2.3.]). There is no difference between the treatment of arrows in T and the rest, except that for $a \in T$, the corresponding band is planar whereas for $a \notin T$ the band goes over the associated 1-handle and it may go over/under other such bands. Note that these connected sum operations are performed directly on the Lagrangian projection without reverting back to the front projection. To justify this, observe that one could use the front projection and perform the connected sum there then one would end up with a more complicated diagram similar to the diagram given in the middle part of Figure 1. However, as in that example, there is a simplifying Reidemeister III move that gives the bottom part of Figure 1. This is a local move and we apply this move in all our connected

sums right away. Figures 5, 6 and 7 describe how the Legendrian L_Γ is constructed in a sufficiently complicated example, from which the general pattern is clear.




FIGURE 6. The configuration of bands and different types of secondary crossings. At the intersection of two bands, the one induced by the arc with the smaller slope in Figure 5 goes over the other. For a more detailed local picture around an unknot see Figure 7.

Now that we have a standard Lagrangian projection of L_Γ , here are the crossings:

- a pair of crossings c_a and c_a^* between $U_{s(a)}$ and $U_{t(a)}$ for each arrow a of Γ
- a self-crossing τ_v of U_v for each vertex v of Γ
- a self-crossing ζ_a of $U_{s(a)}$ near the right foot of the 1-handle h_a , for each arrow a in $\Gamma \setminus T$.
- additional crossings associated to nonplanar bands in the above construction of L_Γ which go over/under each other

3.2. Computation of $\mathcal{B}_\Gamma = CE^*(L_\Gamma)$. The computations in this section are based on the simplification of the Chekanov-Eliashberg DG-algebra of a Legendrian link in $\#^k(S^1 \times S^2, \xi_{std})$ presented in Sec. (2.1), where the internal DG-algebra \mathcal{S}_n is proved to be quasi-isomorphic to its finitely generated DG-subalgebra \mathcal{A}_n as well as the A_∞ -algebra \mathcal{K}_n .

The resolution of L_Γ to be used is the one constructed in Sec. (3.1) with the choice of orientation and base-points indicated in Fig. (7). In this resolution, for every 1-handle h_a induced by an arrow $a \in \Gamma \setminus T$, there are exactly two strands going over h_a which belong to the same component of L_Γ . Therefore the internal generators $c_{12;a}^0, c_{21;a}^1$ can be replaced by the generators of $\mathcal{S}_2^{\mathbb{K}} \simeq \mathbb{K}[z_a, z_a^{-1}]$ with grading 0 and the rest of the internal generators associated to h_a can be dropped from the presentation.

Except for t_v^\pm and z_a^\pm , all the generators of the Chekanov-Eliashberg algebra \mathcal{B}_Γ come from the crossings in the resolution. We use the same notation for these generators as the corresponding crossings, and list them as follows (see Fig. (7)).




FIGURE 7. A local picture showing the choice of attachments and the corresponding computation of the differential.

- a pair of generators c_a, c_a^* with $|c_a| = |c_a^*| = 0$ for each arrow a of Γ ,
- a generator τ_v with $|\tau_v| = -1$ and ²

$$\partial \tau_v = \prod_{t(a)=v} (e_v + c_a c_a^*) - t_v \prod_{s(a)=v, a \notin T} z_a \prod_{s(a)=v, a \in T} (e_v + c_a^* c_a)$$

for each vertex v of Γ , where $s(a)$ and $t(a)$ denote the source and target vertices of a , respectively,

²Although our sign conventions for CE^* follow [12], as we must choose the null-cobordant spin structures on the components of L_Γ , we apply the substitution $t_v \mapsto -t_v$ to their formulas (see [12, Rem2.7])

- a generator ζ_a with $|\zeta_a| = -1$ and

$$\partial\zeta_a = e_v + c_a^* c_a - z_a$$

for each arrow a in $\Gamma \setminus T$, where $v = s(a)$,

- *secondary generators* associated to crossings between nonplanar bands in the construction of L_Γ

We call the generators t_v^\pm , z_a^\pm and those in the first three classes above *primary generators*. In the next lemma, we show that the secondary generators are really an artefact of the projection and could be removed.

Lemma 16. *The DG-subalgebra \mathcal{P}_Γ generated by primary generators of \mathcal{B}_Γ is quasi-isomorphic to \mathcal{B}_Γ .*

Proof. We will prove that \mathcal{B}_Γ and \mathcal{P}_Γ by showing that the secondary generators can be destabilised (after suitable elementary automorphisms) without effecting the rest of \mathcal{B}_Γ . Technically, such a stable tame isomorphism argument requires the DG-algebra to be semi-free, but this does not become an issue in the current set-up since we will only use elementary automorphisms which are identity on the non-free generators t_v^\pm and z_a^\pm .

First of all, note that every intersection point of the arcs in the lower half of the standard form of Γ (blue arcs in Fig. (5)) induces a set of 4 generators which correspond to the crossings between the associated bands in the resolution of L_Γ . We always assume that the band with the smaller slope goes under the other one. A consequence of this is that a polygon whose all vertices are induced by intersection of arcs (in the standard form of Γ) has at least 2 (Reeb-)positive vertices, hence do not contribute to the differential in \mathcal{B}_Γ . This is because at a vertex, the sign (of the quadrant in the interior of the polygon) is negative if and only if the slope decreases while traversing the polygon in the clockwise direction. Moreover, no secondary generator appears in the differential of a secondary generator induced by another intersection or a primary generator.

In the rest of the proof we explain how to destabilise the secondary generators explicitly. The generators induced by the same intersection point in the standard form of Γ are destabilised simultaneously. This process varies slightly depending on whether the intersecting arcs end on the left or on the right of the diagram. Examples of different cases are indicated in Fig. (6). In the arguments below, the arrows in play are $a, a' \in \Gamma \setminus T$ from i, i' to j, j' , resp., and the generators are denoted by p, q, r, s .

Case 1: Both arcs end on the left: The differentials of the generators are

$$\partial p = z_a q + r z_{a'}, \quad \partial q = s z_{a'}, \quad \partial r = -z_a s, \quad \partial s = 0$$

Once we apply the following elementary automorphisms successively

$$s \mapsto z_a^{-1} s, \quad q \mapsto q - z_a^{-1} r z_{a'}, \quad q \mapsto z_a^{-1} q$$

the differentials become

$$\partial p = q, \quad \partial q = 0, \quad \partial r = -s, \quad \partial s = 0$$

and p, q, r, s can be destabilised.

Case 2: Both arcs on the right: In this case, the differentials are

$$\partial p = (e_i + c_a^* c_a)q + r(e_{i'} + c_{a'}^* c_{a'}) , \quad \partial q = s(e_{i'} + c_{a'}^* c_{a'}) , \quad \partial r = -(e_i + c_a^* c_a)s , \quad \partial s = 0$$

We apply the elementary automorphisms

$$p \mapsto p + \zeta_a q + r\zeta_{a'} + \zeta_a s\zeta_{a'} , \quad q \mapsto q + s\zeta_{a'} \quad r \mapsto r - \zeta_a s$$

successively, making the differentials identical to those considered in the first case.

Case 3: The arcs end on different sides: The differentials are

$$\partial p = z_a q + r(e_{i'} + c_{a'}^* c_{a'}) , \quad \partial q = s(e_{i'} + c_{a'}^* c_{a'}) , \quad \partial r = z_a s , \quad \partial s = 0$$

In order to reduce to the first case, it suffices to apply the following elementary automorphisms successively

$$p \mapsto p + r\zeta_{a'} , \quad q \mapsto q + s\zeta_{a'}$$

□

Theorem 17. *Let Γ be a finite graph. The Chekanov-Eliashberg algebra \mathcal{B}_Γ of the Legendrian link L_Γ and the derived multiplicative preprojective algebra \mathcal{L}_Γ associated to Γ are quasi-isomorphic DG-algebras.*

Proof. By the previous lemma, it suffices to find a quasi-isomorphism between \mathcal{P}_Γ and \mathcal{L}_Γ . In fact, there is a DG-algebra isomorphism $\varphi : \mathcal{P}_\Gamma \rightarrow \mathcal{L}_\Gamma$ defined on the primary generators as identity except for

$$\tau_v \mapsto \tau_v + t_v \left(\sum_{i=1}^{\iota_v} z_{a_1} \cdots z_{a_{i-1}} \zeta_{a_i} (e_v + c_{a_{i+1}}^* c_{a_{i+1}}) \cdots (e_v + c_{a_{\iota_v}}^* c_{a_{\iota_v}}) \right) \prod_{s(a)=v, a \in T} (e_v + c_a^* c_a)$$

where $\{a_1, \dots, a_{\iota_v}\}$ is the (ordered) set of arrows with $s(a_i) = v$ and $a_i \in \Gamma \setminus T$. □

4. PLUMBINGS OF COTANGENT BUNDLES OF SURFACES OF POSITIVE GENUS

In this section, we generalise the construction and the computation in the previous section to plumbings of cotangent bundles of surfaces of arbitrary genus.

4.1. A Legendrian surgery presentation. In Fig. (8), we give an example illustrating how to draw a Legendrian surgery picture of a plumbing of cotangent bundles of surfaces of varying genus according to an arbitrary plumbing graph. Inside the boxes, we put a standard surgery picture of the cotangent bundle of a genus g_v surface. This is described in Fig. (10). It is obtained by band summing the 2-handles of g_v copies of the standard picture T^*T^2 given in Fig. (9) and then applying simplifying Reidemeister moves. There is a freedom in choosing the location of the band sum. Our choice makes it easier to see directly that the resulting DG-algebra is quasi-isomorphic to the derived version of the higher genus multiplicative preprojective algebra.





FIGURE 8. Surgery diagram for plumbings of cotangent bundles of surfaces of varying genus. The drawn example shows a plumbing of two copies of T^*T^2 with one copy of T^*S^2 .

4.2. Computation of the Chekanov-Eliashberg DG-algebra. As shown in Lem. (16) the secondary generators induced by intersecting blue curves as in Fig. (6) can be destabilised without effecting the rest of the DG-algebra $\mathcal{B}_{\Gamma, \mathbf{g}}$. In the positive genus case, we have a new set of secondary generators induced by intersecting red and blue curves as in Fig. (8) which can be destabilised exactly as before. Another set of secondary generators $p_{v,k}, q_{v,k}, r_{v,k}$ and $s_{v,k}$, for $k = 1, \dots, g_v - 1$, indicated by quadruples (p_i, q_i, r_i, s_i) for $i = 1, 2, \dots, g - 1$ in Fig. (10), have a configuration and destabilisation similar to those treated in the first case of the proof of Lem. (16). By these initial simplifications, $\mathcal{B}_{\Gamma, \mathbf{g}}$ is quasi-isomorphic to its DG-subalgebra $\mathcal{P}_{\Gamma, \mathbf{g}}$ generated by the primary generators below:

FIGURE 9. Legendrian surgery picture of T^*T^2

- generators t_v^\pm for each vertex v ,
- generators ζ_a, z_a^\pm for each arrow a in $\Gamma \setminus T$,
- generators c_a, c_a^* for each arrow a in Γ ,
- generators $\tau_v, z_{v,i}^\pm, \xi_{v,j}, x_{v,j}$ for each vertex $v, i = 1, \dots, 2g_v$, and $j = 1, \dots, 4g_v$

For the last group of generators above, see Fig. (10) where the notation is simplified by using x_i instead of $x_{v,i}$, etc. Note that $z_{v,i}^\pm$ correspond to the generators of the internal DG-algebra $\mathcal{S}_2^{\mathbb{K}} \simeq \mathbb{K}[z^\pm]$ of the i^{th} 1-handle h_i in the diagram of the cotangent bundle of the surface of genus g_v in Fig. (10).

As in the genus 0 case, the DG-algebra $\mathcal{P}_{\Gamma, \mathbf{g}}$ of primary generators is supported in nonpositive gradings with τ_v, ζ_a and $\xi_{v,j}$ in grading -1 and the rest of the generators in grading 0. Note that although the generators $\xi_{v,4i-2}, \xi_{v,4i-1}$, for $i = 1, \dots, g_v$, are induced by the twists near the right feet of the corresponding 1-handles similar to the generators of type ζ_a , they can be destabilised and will not appear in the ultimate presentation of $\mathcal{B}_{\Gamma, \mathbf{g}}$.

Before we show how to destabilise the generators $\xi_{v,j}$ and $x_{v,j}$, let us describe the differential on $\mathcal{P}_{\Gamma, \mathbf{g}}$ as given by Fig. (10) where $z_i = z_{v,i}, \xi_j = \xi_{v,j}, x_j = x_{v,j}, g = g_v$:

$$\begin{aligned}
 \partial \tau_v &= \prod_{t(a)=v} (e_v + c_a c_a^*) - t_v x_1 \prod_{s(a)=v, a \notin T} z_a \prod_{s(a)=v, a \in T} (e_v + c_a^* c_a) \\
 \partial \zeta_a &= e_{s(a)} + c_a^* c_a - z_a \\
 \partial \xi_{4k-3} &= -z_{2k} x_{4k-3} + x_{4k-2} x_{4k-1} x_{4k}, \text{ for } k = 1, \dots, g-1 \\
 \partial \xi_{4k-2} &= x_{4k-2} - z_{2k-1}, \text{ for } k = 1, \dots, g \\
 \partial \xi_{4k-1} &= x_{4k-1} - z_{2k}, \text{ for } k = 1, \dots, g \\
 \partial \xi_{4k} &= e_v - x_{4k+1} z_{2k-1} x_{4k}, \text{ for } k = 1, \dots, g-2 \\
 \partial \xi_{4g-3} &= -z_{2g} x_{4g-3} + x_{4g-2} x_{4g-1} \\
 \partial \xi_{4g-4} &= e_v - x_{4g-3} x_{4g} z_{2g-3} x_{4g-4} \\
 \partial \xi_{4g} &= e_v - z_{2g-1} x_{4g}
 \end{aligned}$$

The following lemma suffices to generalise the quasi-isomorphism between $\mathcal{B}_{\Gamma, \mathbf{g}}$ and $\mathcal{L}_{\Gamma, \mathbf{g}}$ proved in Thm. (17) for $\mathbf{g} = 0$.




FIGURE 10. The green part of the 2-handle of $T^*\Sigma_g$ is analogous to Fig. (7)

Lemma 18. *The Chekanov-Eliashberg DG-algebra $\mathcal{B}_{\Gamma, \mathbf{g}}$ of the Legendrian link $L_{\Gamma, \mathbf{g}}$ associated to a graph Γ and an s -tuple \mathbf{g} is quasi-isomorphic to the DG-algebra generated by*

- generators τ_v, t_v^\pm for each vertex v ,
- generators ζ_a, z_a^\pm for each arrow a in $\Gamma \setminus T$,
- generators c_a, c_a^* for each arrow a in Γ ,
- generators $\alpha_{v,i}^\pm, \beta_{v,i}^\pm$ for each vertex $v, i = 1, \dots, g_v$

with $|\zeta_a| = |\tau_v| = -1, |t_v^\pm| = |z_a^\pm| = |c_a| = |c_a^*| = |\alpha_{v,i}^\pm| = |\beta_{v,i}^\pm| = 0$ and nontrivial differentials

$$\partial \zeta_a = e_v + c_a^* c_a - z_a$$

$$\partial \tau_v = \prod_{t(a)=v} (e_v + c_a c_a^*) - t_v \prod_{i=1}^{g_v} [\alpha_{v,i}, \beta_{v,i}] \prod_{s(a)=v, a \notin T} z_a \prod_{s(a)=v, a \in T} (e_v + c_a^* c_a)$$

where $[x, y] = xyx^{-1}y^{-1}$.

Proof. We start with the presentation of $\mathcal{P}_{\Gamma, \mathbf{g}} \simeq \mathcal{B}_{\Gamma, \mathbf{g}}$ preceding the statement and (as in the proof of Lem. (16)) utilise an extended notion of stable tame isomorphism where all the elementary automorphisms are identity on the non-free generators $t_v^\pm, z_a^\pm, z_i^\pm$. With the help of such automorphisms, we destabilise the pairs $(\xi_{v,j}, x_{v,j})$ for all j . We do this in groups of four, starting with larger j and moving our way down to $j = 1$. Note that the treatment of the initial group is slightly different than the rest because of the unique feature of the lowest piece of Fig. (10). (In the rest of the proof, we fix v and use the simplified notation g, x_j, ξ_j and z_j for $g_v, x_{v,j}, \xi_{v,j}$, and $z_{v,j}$, respectively.)

The first four automorphisms we apply are

$$x_{4g} \mapsto z_{2g-1}^{-1}(x_{4g} + e_v), \quad x_{4g-1} \mapsto x_{4g-1} + z_{2g}, \quad x_{4g-2} \mapsto x_{4g-2} + z_{2g-1}$$

followed by

$$\xi_{4g-3} \mapsto \xi_{4g-3} + \xi_{4g-2}(x_{4g-1} + z_{2g}) + z_{2g-1}\xi_{4g-1}, \quad \xi_{4g-4} \mapsto \xi_{4g-4} + x_{4g-3}z_{2g-1}^{-1}\xi_{4g}z_{2g-3}x_{4g-4}$$

then

$$x_{4g-3} \mapsto z_{2g}^{-1}(x_{4g-3} + z_{2g-1}z_{2g})$$

and finally

$$\xi_{4g-4} \mapsto \xi_{4g-4} + z_{2g}^{-1}\xi_{4g-3}z_{2g-1}^{-1}z_{2g-3}x_{4g-4}$$

After these automorphisms, the relevant differentials are changed as follows

$$\begin{aligned} \partial \xi_{4g} &= -x_{4g}, \quad \partial \xi_{4g-1} = x_{4g-1}, \quad \partial \xi_{4g-2} = x_{4g-2}, \quad \partial \xi_{4g-3} = -x_{4g-3} \\ \partial \xi_{4g-4} &= e_v - z_{2g}^{-1}z_{2g-1}z_{2g}z_{2g-1}^{-1}z_{2g-3}x_{4g-4} = e_v - [z_{2g}^{-1}, z_{2g-1}]z_{2g-3}x_{4g-4} \end{aligned}$$

and hence we can destabilise the pairs (ξ_i, x_i) for $i = 4g - 3, \dots, 4g$.

The next four automorphisms will help destabilise the next four pairs (ξ_i, x_i) of generators and establish a pattern. These are

$$x_{4g-4} \mapsto z_{2g-3}^{-1}[z_{2g-1}, z_{2g}^{-1}](x_{4g-4} + e_v), \quad x_{4g-5} \mapsto x_{4g-5} + z_{2g-2}, \quad x_{4g-6} \mapsto x_{4g-6} + z_{2g-3}$$

followed by

$$\begin{aligned} \xi_{4g-7} \mapsto & \xi_{4g-7} + (\xi_{4g-6}(x_{4g-5} + z_{2g-2}) + z_{2g-3}\xi_{4g-5}) z_{2g-3}^{-1} [z_{2g-1}, z_{2g}^{-1}] (x_{4g-4} + e_v) \\ & - z_{2g-3} z_{2g-2} z_{2g-3}^{-1} [z_{2g-1}, z_{2g}^{-1}] \xi_{4g-4} \end{aligned}$$

then

$$x_{4g-7} \mapsto z_{2g-2}^{-1} (x_{4g-7} + z_{2g-3} z_{2g-2} z_{2g-3}^{-1} [z_{2g-1}, z_{2g}^{-1}])$$

and

$$\xi_{4g-8} \mapsto \xi_{4g-8} + z_{2g-2}^{-1} \xi_{4g-7} z_{2g-5} x_{4g-8}$$

These automorphisms result in the following differentials

$$\begin{aligned} \partial \xi_{4g-4} &= -x_{4g-4}, \quad \partial \xi_{4g-5} = x_{4g-5}, \quad \partial \xi_{4g-6} = x_{4g-6}, \quad \partial \xi_{4g-7} = -x_{4g-7} \\ \partial \xi_{4g-8} &= e_v - [z_{2g-2}^{-1}, z_{2g-3}] [z_{2g-1}, z_{2g}^{-1}] z_{2g-5} x_{4g-8} \end{aligned}$$

Applying $(g-2)$ additional sets of four automorphisms analogous to the last four above, we destabilise all the pairs (ξ_i, x_i) . This turns the differential of τ_v into

$$\partial \tau_v = \prod_{t(a)=v} (e_v + c_a c_a^*) - t_v P_C \prod_{s(a)=v, a \notin T} z_a \prod_{s(a)=v, a \in T} (e_v + c_a^* c_a)$$

where P_C is the product of commutators

$$\begin{aligned} & [z_2^{-1}, z_1] [z_6^{-1}, z_5] \cdots [z_{2g-2}^{-1}, z_{2g-3}] [z_{2g-1}, z_{2g}^{-1}] \cdots [z_7, z_8^{-1}] [z_3, z_4^{-1}] \text{ if } g \text{ is even, and} \\ & [z_2^{-1}, z_1] [z_6^{-1}, z_5] \cdots [z_{2g}^{-1}, z_{2g-1}] [z_{2g-3}, z_{2g-2}^{-1}] \cdots [z_7, z_8^{-1}] [z_3, z_4^{-1}] \text{ if } g \text{ is odd} \end{aligned}$$

In order to match this last differential with the expression in the statement (which makes it easier to see the quasi-isomorphism $\mathcal{B}_{\Gamma, \mathbf{g}} \simeq \mathcal{L}_{\Gamma, \mathbf{g}}$), we simply relabel the generators z_i^\pm so that

$$\alpha_{v,i} = z_{4i-2}^{-1}, \quad \alpha_{v,j} = z_{4g-4j+3}, \quad \beta_{v,i} = z_{4i-3}, \quad \beta_{v,j} = z_{4g-4j+4}, \quad \text{for } 1 \leq i \leq \frac{g+1}{2} < j \leq g$$

□

Proof of Thm. (1): With the above lemma established, we proceed as in the proof of Thm. (17) and consider the following isomorphism $\varphi : \mathcal{P}_{\Gamma, \mathbf{g}} \rightarrow \mathcal{L}_{\Gamma, \mathbf{g}}$ of DG-algebras defined to be the identity on all primary generators except for

$$\tau_v \mapsto \tau_v + t_v \prod_{i=1}^{g_v} [\alpha_{v,i}, \beta_{v,i}] \left(\sum_{i=1}^{t_v} z_{a_1} \cdots z_{a_{i-1}} \zeta_{a_i} (e_v + c_{a_{i+1}} c_{a_{i+1}}^*) \cdots (e_v + c_{a_{t_v}} c_{a_{t_v}}^*) \right) \prod_{s(a)=v, a \in T} (e_v + c_a^* c_a)$$

where $\{a_1, \dots, a_{t_v}\}$ is the (ordered) set of arrows with $s(a_i) = v$ and $a_i \in \Gamma \setminus T$. □

Remark 19. Throughout the paper, we only considered plumbings with positive intersections. In the case of negative plumbings, the Chekanov-Eliashberg DG-algebra of the corresponding Legendrian link can be described by the same set of generators as above but the differential gets modified as follows

$$\partial \tau_v = \prod_{t(a)=v, \text{sgn}(a)=+} (e_v + c_a c_a^*) - t_v \prod_{i=1}^{g_v} [\alpha_{v,i}, \beta_{v,i}] \prod_{t(a)=v, \text{sgn}(a)=-} (e_v + c_a c_a^*) \prod_{s(a)=v} (e_v + c_a^* c_a),$$

where $\text{sgn}(a)$ denotes the sign of the plumbing associated to the arrow a . Importantly, the DG-algebra is no longer non-positively graded since c_a and c_a^* have gradings ± 1 whenever $\text{sgn}(a) = -$.

REFERENCES

- [1] M. Abouzaid, A geometric criterion for generating the Fukaya category. *Publ. Math. Inst. Hautes Études Sci.* 112 (2010), 191–240.
- [2] M. Abouzaid, I. Smith, Exact Lagrangians in plumbings. *Geom. Funct. Anal.* 22 (2012), no. 4, 785–831.
- [3] R. Bezrukavnikov, M. Kapranov, Microlocal sheaves and quiver varieties. *Ann. Fac. Sci. Toulouse Math.* (6) 25 (2016), no. 2-3, 473–516.
- [4] P. Boalch, Global Weyl groups and a new theory of multiplicative quiver varieties. *Geom. Topol.* 19 (2015), no. 6, 3467–3536.
- [5] F. Bourgeois, T. Ekholm, Y. Eliashberg, Effect of Legendrian surgery. With an appendix by S. Ganatra and M. Maydanskiy, *Geom. Topol.* 16 (2012), no. 1, 301–389.
- [6] R. Casals, E. Murphy, Legendrian Fronts for Affine Varieties. Preprint, arXiv:1610.0697,
- [7] Y. Chekanov, Differential algebra of Legendrian links. *Invent. Math.* 150 (2002), no. 3, 441–483.
- [8] W. Crawley-Boevey, P. Shaw, Multiplicative preprojective algebras, middle convolution and the Deligne-Simpson problem. *Adv. Math.* 201 (2006), no. 1, 180–208.
- [9] B. Chantraine, G. Dimitroglou Rizell, P. Ghiggini, R. Golovko Geometric generation of the wrapped Fukaya category of Weinstein manifolds and sectors, preprint arXiv:1712.09126
- [10] V. Drinfeld, DG quotients of DG categories. *J. Algebra* 272 (2004), no. 2, 643–691.
- [11] T. Ekholm, Y. Lekili, Duality between Lagrangian and Legendrian invariants. Preprint, arXiv:1701.01284.
- [12] T. Ekholm, L. Ng, Legendrian contact homology in the boundary of a subcritical Weinstein 4-manifold. *J. Differential Geom.* 101 (2015), no. 1, 67–157.
- [13] Y. Eliashberg, Invariants in contact topology. *Proceedings of the International Congress of Mathematicians, Vol. II (Berlin, 1998). Doc. Math.* 1998, Extra Vol. II, 327–338.
- [14] T. Etinger, Y. Lekili, Koszul duality patterns in Floer theory. *Geom. Topol.* 21 (2017), 3313–3389.
- [15] R. Gompf, Handlebody construction of Stein surfaces. *Ann. of Math.* (2) 148 (1998), no. 2, 619–693.
- [16] F. Haiden, L. Katzarkov, M. Kontsevich, Flat surfaces and stability structures. *Publ. Math. Inst. Hautes Études Sci.* 126 (2017), 247–318.
- [17] A. Keating, Lagrangian tori in four-dimensional Milnor fibres. *Geom. Funct. Anal.* 25 (2015), no. 6, 1822–1901.
- [18] J.-I. Miyachi, Localization of Triangulated Categories and Derived Categories. *Jour. of Algebra.* 141 (1991), 463–483.
- [19] D. Nadler, E. Zaslow, Constructible sheaves and the Fukaya category. *J. Amer. Math. Soc.* 22 (2009), no. 1, 233–286.
- [20] L. Ng, Computable Legendrian invariants. *Topology* 42 (2003), no. 1, 55–82.
- [21] T. Pantev, B. Toën, M. Vaquié, G. Vezzosi, Shifted symplectic structures. *Publ. Math. Inst. Hautes Études Sci.* 117 (2013), 271–328.
- [22] J. Pascaleff, Floer cohomology in the mirror of the projective plane and a binodal cubic curve. *Duke Math. J.* 163 (2014), no. 13, 2427–2516.
- [23] P. Seidel, Graded Lagrangian submanifolds. *Bull. Soc. Math. France* 128 (2000), no. 1, 103–149.
- [24] P. Seidel, Fukaya categories and Picard-Lefschetz theory. *Zurich Lectures in Advanced Mathematics*, European Mathematical Society (EMS), Zürich, 2008.
- [25] V. Shende, A. Takeda, Symplectic structures from topological Fukaya categories, Preprint, arXiv:1605.02721.

- [26] D. Yamakawa, Geometry of multiplicative preprojective algebra. Int. Math. Res. Pap. IMRP 2008, Art. ID rpn008, 77pp.
- [27] W-K Yeung, Relative Calabi-Yau completions. Preprint, arXiv:1612.06352.

DEPARTMENT OF MATHEMATICS, KOÇ UNIVERSITY, SARIYER, ISTANBUL 34450 TURKEY
Email address: `tetgu@ku.edu.tr`

KING'S COLLEGE LONDON, STRAND, LONDON, UK
Email address: `yanki.lekili@kcl.ac.uk`